

27-28.

DISSERTATIO ENTOMOLOGICA,  
INSECTA FENNICA  
ENUMERANS,

---

CUJUS PARTICULAM DECIMAM QUINTAM,  
CONS. AMPL. FACULT. PHILOS. ABOËNSIS,

PRÆSIDE

CAROLO REGINALDO SAHLBERG,

*Phil. & Med. Doct. Oecon. & Hist. Natur. Prof.  
Publ. & Ord. Collegii Imp. Med. Assess. Ord. Imp.  
de S. Wolodimiro in quart. Class. Equ. Societ. Imp.  
Nat. Studios. Moscov. Societ. Nat. Scrut. Lips. &  
Physiogr. Lund. nec non Soc. Imp. Oecon.  
Fenn. Membro,*

PUBLICÆ CENSURÆ OFFERT

PRO LAUREA

PETRUS PHILIPPUS LINDFORSS,  
Stip. Publ. Nyland.

In Auditorio Medico die XXVI Maji 1827.

h. a. m. c.

---

ABOË, Typis FRENCKELLIORUM,

UNIVERSITY OF CHICAGO

PHYSICS DEPARTMENT

PHYSICS 551

PROFESSOR

PHYSICS

PHYSICS

PHYSICS

PHYSICS

PHYSICS

PHYSICS

PHYSICS

*Gyll. Ins. Svec. 2. 77. 1.*

*Taphria vivalis. Dej. Catal. Coleopt. p. 10.*

*Carabus rotundatus var. b. Schönh. Synon. 1. 214, 258. β. Payk. Faun. Svec. 1. 136. 50. var β.*

Habitat in truncis putridis & sub lapidibus minus frequens.

Obs. Variat corpore toto plus minus rufo-testaceo. Forte nuper exclusus?

## PANAGÆUS.

*Palpi medii & postici articulo ultimo crassiore, obconico, oblique truncato, subsecuriformi. Mandibula brevissima valde arcuata, interne integra. Labium dente intermedio bifido. Maxilla arcuata acutissima, intus ciliata. Caput postice in collum coarctatum. Thorax rotundatus.*

1. *P. Crux*: pubescens ater, elytris punctato-striatis rufis, basi, cruce communi apiceque nigris. Long. 3 lin.

*Sturm Ins. Germ. 3. 170. 1. Dej. Spec. Coleopt. 2. 286. 3.*

*Carabus Crux major. Schönh. Synon. 1. 209. 238. Payk. Faun. Svec. 1. 137 52.*

Habitat sub lapidibus in Fennia rarissime. In Tavastia; *D. Boucht*; in Parœcia Padasjoki *D. v. Nordtman*.

## HARPALUS.

*Palpi medii & postici articulo ultimo elongato cylindrico, apice truncato, interdum*

*attenuato subacuminato. Mandibula arcuata, acuta, inermis. Ligula elongata, trifida, lacinia media coriacea, apice obtusa vel truncata, lateralibus membranaceis. Caput elongatum, anterius angustatum. Elytra longitudine abdominis.*

Familia 1. Thorace oblongo postice angustiore subcordato, basi truncato, angulis acutis.

Sect. 1. Alati.

1. *H. leucophthalmus*: nigro-piceus lævis subdepressus, elytris subtiliter punctato-striatis, femorum posticorum stipulis acuminatis. Long. 8 lin.

*Gyll. Ins. Svec. 2. 80. 1.*

*Carabus planus. Schönh. Synon. 1. 180. 63.*

*Carabus spiniger. Payk. 1. 114. 23.*

*Sphodrus planus. Sturm Ins. Germ. 5. 149. 1.*  
*Dej. Catal. Coleopt. p. 10.*

Habitat sub lapidibus & foliis deciduis, locis umbrosis, rarius.

2. *H. angusticollis*: niger nitidus subdepressus, thorace angusto postice coarctato late marginato, angulis posticis parum prominulis acutis fere re-ctis, elytris striatis, apice sinuato-emarginatis. Long. 4 lin.

*Gyll. Ins. Svec. 2. 81. 2.*

*Carabus angusticollis. Schönh. Synon. 1.*  
*185. 88.*

*Platynus angusticollis*. Dej. *Catal. Coleopt.*  
p. 10.

*Anchomenus angusticollis*. Sturm *Ins. Germ.*  
5. 168. 2.

*Carabus assimilis*. Payk. *Faun. 1. 119. 30.*

Var. b. Supra nigro-piceus, subtus pallidior.

Habitat sub lapidibus, locis humidis, passim.

Var. b. ad Wasam; D. *Wasastjerna.*

3. *H. albipes*: oblongus nigro-piceus, antennis, palpibus, pedibus elytrorumque limbo pallidis, thorace ad basin punctato, elytris leviter striatis. Long. 3 lin.

*Gyll. Ins. Svec. 2. 82. 3.*

*Anchomenus albipes*. Sturm *Ins. Germ. 5.*  
175. 6.

*Anchomenus pallipes*. Dej. *Catal. Coleopt.*  
p. 10.

*Carabus albipes*. Schönh. *Synon. 1. 190. 116.*  
Payk. *Faun. Svec. 3. App. 442. 35—36.*

Habitat sub lapidibus, in Fennia Australi, rarius; in Hortis Aboënsibus interdum haud infrequens.

4. *H. prasinus*: oblongus niger, palpis antennarum basi pedibusque testaceis, capite thoraceque viridi-æneis, elytris ferrugineis, macula communi postica cyanea. Long. 2½ lin.

*Gyll. Ins. Svec. 2. 83. 4.*

*Anchomenus prasinus*, Dej. Catal. Coleopt. p. 10. — Sturm Ins. Germ. 5. 171. 4.

*Carabus prasinus*, Schönh. Synon. 1. 215. 264. Payk. Faun. Svec. 1. 126. 38.

Habitat ad Wasam rarissime. D, Wasastjerna.

5. *H. oblongo-punctatus*: nigro-æneus, palpis tibiis tarsisque rufescentibus, elytris leviter striatis, foveolis pluribus serie irregulari impressis. Long. 4 lin.

Gyll. Ins. Svec. 2. 85. 6.

*Pterostichus oblongo-punctatus*, Dej. Catal. Coleopt. p. 12.

*Platysma oblongo-punctata*, Sturm Ins. Germ. 5. 51. 8.

*Carabus oblongo-punctatus*, Schönh. Synon. 1. 186. 92. Payk. Faun. Svec. 1. 121. 32.

Var b. niger, palpis tibiis tarsisque nigro-piceis.

Habitat sub muscis, lapidibus, truncis, locis sylvaticis, frequens. Var. b. in Lapponia. D, Kolström,

6. *H. niger*: subdepressus niger, thorace postice utrinque bistriato, elytris profunde striatis, interstitiis convexis. Long. 6—7 lin.

Gyll. Ins. Svec. 2. 86. 7.

*Platysma nigra*, Dej. Catal. Coleopt. p. 12.

*Pterostichus niger*, Sturm Ins. Germ. 5. 5. 1.

*Carabus niger*, Schönh. Synon. 1. 179. 62.

*Carabus striatus*. Payk. Faun. Svec. 1. 115. 24.

Habitat sub lapidibus & in truncis putridis sat frequens.

7. H. Nigrita: niger nitidus immaculatus, thorace brevior, lateribus æqualiter rotundato, postice parum angustiore, utrinque fovea magna punctata impressa, elytrorum striis sublævibus. Long. lin. 4.

Gyll. Ins. Svec. 2. 88. 8.

*Omasus nigrita*. Dej. Catal. Coleopt. p. 12.

*Platysma nigrita*. Sturm Ins. Germ. 5. 64. 15.

*Carabus Nigrita*. Schönh. Synon. 1. 208. 223.

Payk. Faun. Svec. 1. 157. 77.

Var. b. Ferrugineo-testaceus, capite thoracisque disco nigro-brunneis.

Gyll. Ins. Svec. 1. c. var b.

Habitat sub muscis & lapidibus, locis humidis, frequens. Var. b mihi numquam obvia.

8. H. minor: \*) oblongus niger nitidus, antennis pedibusque rufo-piceis, thorace angustiore utrinque bistriato, striis elytrorum subcrenatis. Long.  $2\frac{3}{4}$  lin.

*H. anthracinus*. Gyll. Ins. Svec. 2. 89. 9.

*Omasus minor*. Dej. Catal. Coleopt. p. 12.

---

\*) Loco nominis specifici, *anthracinus*, speciei ab hac diversæ antea impositi, nomen ab Illustrissimo D. Comite DEJEAN propositum adoptare, necessarium duxi.

*Platysma longicollis*. Sturm Ins. Germ. 5.  
80. 25.

*Carabus Nigrita*. Payk. Faun. Svec. 1. 157. 77.  
var. minor.

Var. b. Ferrugineo-testaceus, subtus dilutior.

Gyll. Ins. Svec. 1. c. var. b.

Habitat sub muscis & lapidibus, locis, humidis  
minus frequens.

9. *H. vernalis*: brevior niger nitidus depressus,  
antennis pedibusque rufo-piceis, thorace latiore  
postice impresso punctulato, utrinque bistriato.  
Long.  $2\frac{1}{2}$  lin.

Gyll. Ins. Svec. 2. 90. 10.

*Argutor vernalis*. Dej. Catal. Coleopt. p. 11.

*Platysma vernalis*. Sturm Ins. Germ. 5.  
69. 18.

*Carabus vernalis*. Schönh. Synon. 1. 217. 274.

Habitat sub muscis & lapidibus, locis humidis,  
passim.

#### Sect. 2 APTERI.

10. *H. subcyaneus*: oblongus nigro-piceus sub-  
depressus, elytris obscure cyaneis, subtiliter pun-  
ctato-striatis, thorace postice profunde impresso.  
Long. 5 lin.

Gyll. Ins. Svec. 2. 91. 11.

*Sphodrus subcyaneus*. Sturm Ins. Germ. 5.  
151. 2.



*Sphodrus Terricola*. Dejean Catal. Coleopt.  
p. 10.

*Carabus subcyaneus*. Schönh. Synon. 1. 183. 78.  
*C. Terricola*. Payk. Faun. 1. 109. 16.

Habitat in Fennia Australi, rarissime. Duo specimenina prope Aboam ceperunt D. Dammert & Tams.

11. *fervus*: oblongus niger thoracis foveis posticis rugoso-punctatis bistriatis, elytris profunde striatis, striis tenue punctulatis. Long. 6 lin.

Habitat in Fennia Australi rarissime.

Descr. *Platysma nigerrima* Sturmii valde similis videtur, sed differt striis elytrorum subtiliter punctulatis. Caput porrectum nigrum nitidum, fovea inter antennis oblonga utrinque. Palpi picei. Antennæ thorace vix longiores, basi nigrae, extrorsum fusco-pubescentes subtus piceæ. Thorax antice longitudine latior, submarginatus, angulis deflexis obtusis, lateribus late marginatus rotundatus, posterius angustatus, basi in medio nonnihil emarginatus, angulis acuminatis prominulis deflexis, supra modice convexus, totus niger nitidus, in medio canaliculatus atque intra basin utrinque fovea magna rugoso-punctata, striis duabus paullo profundioribus exarata, plicaque elevata recta extrorsum terminata, impressus. Scutellum parvum triangulare nigrum nitidum. Elytra thoracis basi paullo latiora, ultra medium parum dilatata, apice sinuato-rotundata, supra parum convexa, nigra nitida, profunde striata, striis tenue punctatis; juxta striam a sutura secundam puncta tria profundius impressa, & in stria a margine secunda series e foveolis, in medio parcius exsculptis. Corpus subtus nigrum nitidum, lateribus thoracis pectorisque punctulatis, segmentis abdominis impressione utrinque sat profunda notatis. Pedes nigri nitidi, spinulis tarsorum piceo-rufis.

12. *H. ater*: oblongus niger immaculatus, thorace foveis posticis bilobis, elytris tenue striatis, striis subtiliter punctulatis. Long. 5 lin.

Habitat in Fennia Australi, rarissime.

Descr. *H. melanari* simillimus & differt tantum striis elytrorum levius impressis & punctulatis. Caput cum palpis & antennis, thorax & scutellum omnino ut in *H. melanari*. Elytra thoracis basi latiora, ultra medium dilatata, apice sinuato-rotundata, supra modice convexa nigra nitida striata, striis leviter impressis & subtiliter punctulatis, juxta striam a sutura secundam puncta duo profundius impressa & series foveolarum in stria a margine secunda ut in præcedente. Corpus & pedes ut in *H. melanari*.

13. *H. melanaris*: oblongus niger immaculatus, thoracis foveis posticis bilobis, elytrorum striis profundis lævibus. Long. 5 lin.

*Gyll. Ins. Svec. 2. 92. 12.*

*Omaseus melanarius, Dej. Catal. Coleopt. p. 12.*

*Platysma leucophthalma, Sturm Ins. Germ. 5. 39. 1.*

*Carabus leucophthalmus, Schönh. Synon. 1. 178. 52. Payk. Faun. Svec. 1. 108. 15.*

Habitat in terra & sub lapidibus, frequens.

14. *H. lepidus*: oblongus supra viridi-cupreus, antennis pedibusque nigris, thorace postice utrinque profunde bistriato, elytris disco tripunctatis. Long.  $4\frac{1}{2}$  lin.

Mas: elytris nitidis, superficie vix coriacea.

Femina: elytris opacis, sericeo-micantibus, superficie evidentius granulato-coriacea.

*Gyll. Ins. Svec. 2. 94. 14.*

*Poecilus lepidus, Dej. Catal. Coleopt. p. 11.*

*Platysma lepida, Sturm Ins. Germ. 5. 92. 33.*

*Carabus lepidus*, Schönh. *Synon.* 1, 194, 151.  
Payk. *Faun. Svec.* 1, 110, 17.

Var. b. Supra cupreus.

*Gyll. Ins. Svec.* 1, c. var. b.

Var. c. Supra viridi-cupreus, elytrorum limbo virescenti-azureo.

*Gyll. Ins. Svec.* 1, c. var. c.

Var. d. Supra viridis.

*Gyll. Ins. Svec.* 1, c. var. d. Schönh. *Synon.* 1, c. var.  $\beta$ .

Var. e. Supra nigro-virescens sive olivaceus.

*Gyll. Ins. Svec.* 1, c. var. e. Schönh. *Synon.* 1, c. var.  $\gamma$ .

Var. f. Supra violaceus.

*Gyll. Ins. Svec.* 1, c. var. f. Schönh. *Synon.* 1, c. var.  $\delta$ .

Var. g. Supra niger.

*Gyll. Ins. Svec.* 1, c. var. g.

Var. h. Capite thoraceque cæruleo-violaceis, elytris nigris, margine violaceo.

*Gyll. Ins. Svec. P. 4. add.*

Var. i. Capite thoraceque viridibus, elytris nigris, margine viridi.

*Gyll. Ins. Svec. P. 4. add.*

Habitat in Fennia Australi minus frequens. In Territorio Wiburgensi a *D. Mannerheim* pluries lectus.

15. *H. arcticus*: oblongus supra cupreus nitidus, subtus cupreo-violaceus, capite thoraceque postice utrinque profunde foveolato crebre punctatis, elytris tenue striato-punctatis, foveolisque dissi subtribus impressis, tibiis ferrugineis. Long.  $3\frac{1}{4}$  lin.

*Gyll. Ins. Svec. 2. 96. 15.*

*Blethisa arctica. Gyll. Ins. Svec. P. 4. add.*

Var. b. Supra virescenti-violaceus.

Habitat in Lapponia rarissime. D. *Grape*. Ad Muonioniska nonnulla specimina legit D. M. *Kolström*.

16. *H. rufipes*: oblongus nigro-piceus, subtus dilutior, pedibus rufis, thorace postice profunde foveolato, elytris punctato-striatis. Long.  $3\frac{1}{4}$  lin.

*Gyll. Ins. Svec. 2. 97. 16.*

*Patrobus rufipes. Dej. Catal. Coleopt. p. 10.*

*Platysma rufipes. Sturm Ins. Germ. 5. 56. 10.*

*Carabus rufipes. Schönh. Synon. 1. 188. 105.*

*Carabus excavatus. Payk. Faun. Svec. 1. 123. 34.*

Var. b. Corpore toto rufo-ferrugineo, capite thoraceque interdum obscurioribus.

*Gyll. Ins. Svec. & Sch. Synon. 1. c. var. b.*

Var. c. Alatus.

*Gyll. Ins. Svec. 1. c. var. c. & 3. 693. Add. 16. c.*

*Patrobus Septentrionalis. Schönh. in litt.*

Var. d. Paulo major, femoribus & interdum antennarum basi nigro-piceis.

*Gyll. Ins. Svec. P. 4. add.*

Habitat in terra & sub lapidibus, locis humidis, passim. Var. c. in Lapponia non infrequens.

17. *H. strenuus*: niger nitidus, antennis pedibusque rufo-ferrugineis, thorace lateribus rotundato, postice angustiore utrinque unistriato, subtus punctato, elytrorum striis evidenter punctatis. Long.  $2\frac{1}{4}$  lin.

*Gyll. Ins. Svec. 2. 98. 17.*

*Argutor strenuus* (*Gyll.*) *Dej. Catal. Coleopt. p. 11.*

*P. atysma strenua*, *Sturm Ins. Germ. 5. 71. 19.*

*Carabus strenuus*, *Schönh. Synon. 1. 179. 60.*

Var. b. Ferrugineus vel fusco-testaceus totus.

*Gyll. Ins. Svec. 1. c. var. b.*

Var. c. Alatus.

*Gyll. Ins. Svec. 1. c. var. c.*

Habitat sub muscis & lapidibus, locis humidis, passim.

18. *H. pullus*: niger nitidus, antennis pedibusque ferrugineis, thorace rotundato postice angustiore, utrinque unistriato, subtus impunctato, elytrorum striis leviter crenatis. Long.  $2\frac{1}{4}$  lin.

*Schönh. in litt.*

*H. eruditus?* Gyll. *Ins. Svec. P. 4. add.*

Var. b. Femoribus piceis.

Gyll. *Ins. Svec. 1. c. var. b.*

Var. c. Supra ferrugineus.

Gyll. *Ins. Svec. 1. c. var. c.*

Habitat sub lapidibus & muscis passim.

19. *H. oblongus*: piceus nitidus, thorace elongato angusto, elytris crenato-striatis fusco-testaceis, antennis pedibusque pallidis. Long.  $2\frac{1}{2}$  lin.

Gyll. *Ins. Svec. 2. 99. 18.*

*Anchomenus oblongus*. Dej. *Catal. Coleopt. p. 10. Sturm Ins. Germ. 5. 173. 5.*

*Carabus oblongus*. Schönh. *Synon. 1. 190. 115.*

*Carabus tæniatus*. Payk. *Faun. Svec. 1. 113. 21.*

Var. b. Rufo-ferrugineus, capite supra nigropiceo, elytris pedibusque pallidis.

Gyll. *Ins. Svec. 1. c. var. b.*

Habitat in Fennia Australi, rarissime; in Yläne semel captus. Var. b. in Fennia nondum inventa.

Fam. 2. Thorace brevi transverso, postice angustiori, angulis acutis reflexis; alati.

20. *H. aulicus*: brevior latus nigropiceus, subtus dilutior, ore, antennis pedibusque rufo-ferrugineis, thorace postice depresso utrinque bistriato, antice posticeque punctato. Long.  $4\frac{1}{2}$  lin.

*Gyll. Ins. Svec. 2. 101. 19.*

*Amara aulica. Dej. Catal. Coleopt. p. 9.*

*Amara picea. Sturm Ins. Germ. 6. 10. 1.*

*Carabus aulicus. Schönh. Synon. 1. 181. 69.*

*Carabus bicolor. Payk. Faun. Sv. 1. 159. 79.*

Habitat in agris, pratis, sub lapidibus, frequens.

21. *H. torridus*: nigro-piceus, antennis palpisque rufo-ferrugineis, thorace postice angustiore, foveis bilobis, elytris punctato-striatis, pedibus rufo-piceis. Long.  $3\frac{1}{2}$  lin.

*Gyll. Ins. Svec. 2. 102. 20.*

*Carabus alpinus* var.  $\beta$ .  $\gamma$ . *Schönh. Synon.*

1. 202. 192.  $\beta$ .  $\gamma$ . *Payk. Faun. Svec. 1. 160.*

80.  $\beta$ . ♂  $\gamma$ .

Ver. b. Corpore subtus, pedibus, labro oreque toto testaceis. Forte immaturus.

*Gyll. Ins. Svec. P. 4. add.*

Habitat in Lapponia, sub lapidibus, haud infrequens.

22. *H. alpinus*: niger, antennarum basi, elytrorum disco pedibusque rufo-ferrugineis, thorace postice parum angustato, foveis bilobis, elytris punctato-striatis. Long.  $3\frac{1}{2}$  lin.

*Gyll. Ins. Svec. 2. 103. 21.*

*Amara alpina. Sturm Ins. Germ. 6. 12. 2.*

*Carabus alpinus. Schönh. Synon. 1. 202. 192.*

*Payk. Faun. Svec. 1. 160. 80.*

Var. b. Capite, thorace elytrorumque limbo viridi-æneis.

*Gyll. Ins. Svec. P. 4. add. var. b.*

Var. c. Nigro-piceus, elytris totis concoloribus.

*Gyll. Ins. Svec. 1. c. var. c.*

Var. d. Nigro-æneus, elytris totis concoloribus.

*Gyll. Ins. Svec. 1. c. var. d.*

Var. e. Viridi-æneus, elytris totis concoloribus.

*Gyll. Ins. Svec. 1. c. var. e.*

Habitat in alpihus Lapponicis, sub lapidibus, passim.

23. *H. apricarius*; ovato-oblongus convexus supra fusco-æneus, thorace postice parum angustato, utrinque bistriato punctato, elytris crenato-striatis. Long. 3 lin.

*Gyll. Ins. Svec. 2. 104. 27.*

*Amara apricaria. Dej. Catal. Coleopt. p. 9.*

*Sturm Ins. Germ. 6. 19. 6.*

*Carabus apricarius. Schönh. Synon. 1. 214. 261. Payk. Faun. Svec. 1. 162. 82.*

Var. b. Supra rufo-piceus, subtus dilutior.

*Gyll. Ins. Svec. 1. c. var. b.*

Habitat sub lapidibus, locis aridis, & ad agrorum margines ubique frequens.

24. *H. fulvus*: brevis latus ferrugineus, thorace postice utrinque bistriato, elytris æneo-micantibus, tenue punctato-striatis. Long.  $3\frac{1}{4}$  lin.

*Gyll. Ins. Svec. 2. 105. 23.*



*Amara fulva*. Dej. Catal. Coleopt. p. 9.

Sturm Ins. Germ. 6. 17. 5.

*Carabus fulvus*. Schönh. Synon. 1. 214. 262.

*Carabus ferrugineus*. Payk. Faun. Svec. 1.  
161. 81.

Var. b. Totus pallidus, oculis tantum obscurioribus.

Gyll. Ins. Svec. 1. c. var. b.

Habitat in terra & sub lapidibus frequens.

25. *H. ruficornis*: oblongus nigro-piceus, antennis pedibusque rufis, elytris striatis pubescentibus opacis, interstitiis crebre punctulatis. Long. 5 lin.

Sturm. Ins. Germ. 4. 8. 2. Dej. Catal. Coleopt. p. 14.

*Carabus ruficornis*. Schönh. Synon. 1. 181. 71.  
Payk. Faun. Svec. 1. 158. 78.

Var. b. Elytris fuscis, corpore subtus testaceo.

Gyll. Ins. Svec. 2. 107. 24. var. b.

Var. c. Duplo minor, subtus rufo-piceus vel testaceus.

Gyll. Ins. Sv. 1. c. var. c. Sturm Ins. Germ. 1. c. var. b.

*H. griseus*. Dej. Catal. Coleopt. p. 14.

*Carabus griseus*. Panz. Faun. Germ. Tab. 38. N:o 1.

Var. d. Supra ferrugineus, subtus pallidus.

Gyll. Ins. Svec. 1. c. var. d.

Habitat sub lapidibus & in terra, frequens. Variet. c. & d. rarius occurrunt.

26. *H. puncticollis*: niger subpubescens, subtus rufo-piceus, antennis pedibusque pallide rufis, thorace profunde punctato, elytris striatis, interstitiis punctulatis. Long. 3 lin.

*Sturm Ins. Germ. 4. 103. 59.*

*Ophonus puncticollis. Dej. Catal. Coleopt. p. 13.*

*Carabus puncticollis. Schönh. Synon. 1. 188. 102. — Payk. Faun. Svec. 1. 120. 31.*

Habitat in Fennia Australi sub lapidibus, locis aridis, minus frequens.

Obs. Thorax in speciminibus Fennicis paullo brevior & parcius punctatus, quam in Svecicis, esse videtur. An diversa species?

27. *H. Forsströmii*: nigro-piceus nitidus punctatus, antennis pedibusque fusco-ferrugineis, thorace, late marginato, elytris punctato-striatis, subacuminatis.

*Gyll. Ins. Svec. 2. 111. 27.*

*Pteroloma Forsströmii. Gyll. Ins. Sv. P. 4 add.*

Habitat in Lapponia rarissime. *D. Forsström.*

Fam. 3. Thorace postice paullo angustiori, angulis aut rectis aut subobtusis, vix prominulis, nec exacte rotundatis.

28. *H. holosericeus*: niger, capite æneo-nitente, thorace rugoso, elytris fuscis, sericeo-pubescentibus striatis, interstitiis rugoso scabris. Long.  $4\frac{1}{2}$  lin.