

PERICULUM ENTOMOGRAPHICUM,
SPECIES INSECTORUM
NON DUM DESCRIPTAS
PROPONENS,

QUOD
VENIA FACULT. PHILOS. ABOËNSIS AMPLISS.

P R A E S I D E
CAROLO REGINALDO SAHLBERG,
*Phil. & Med. Doct., Oecon. & Hist. Nat. Profess.
Publ. & Ord., Collegii Imp. Med. Assess., Societt. Imp.
Nat. Studios. Moscov. & Nat. Scrut. Lips. nec non
Soc. Imp. Oecon. Fenn. Membro,*

P U B L I C Æ C E N S U R Æ S U B M I T T I T

P R O L A U R E A
C A R O L U S T E N G S T R Ö M ,
Ostrobothn.

I n Auditorio Philos. die XVI Junii 1813,

h. a. m. c.

P. V.

A B O Æ , Typis FRENCKELLIORUM.

PERICIA IN INSISTE
ИСЯТОЕ ИНСИСТЕ

ЗАЩИРЕННОМО

ДОЛГОВОДНОСТЬ

Elytra thorace paullo latiora sed quadruplo fere longiora, ad humeros parum prominulos & apice rotundata, supra convexa, lateribus anguste marginata, atra, nitidissima, subtiliter crebreque punctulata.

Corpus subtus pallide testaceum, nitidum, punctulatum, tenuē pubescens.

Pedes pallide testacei, nitidi, punctulati, tenuē pubescentes; tibiarum apex tarsique infuscata.

Var. b. differt elytris basi tantum, ad quartam longitudinis partem, atris, colore thoracis elytrorumque magis rufo-testaceo, atque antennarum apicē tarsisque minus infuscatis.

Var. c. Pallide testacea tota, tarsi tantum infuscata.

44. CRIOCERIS APICALIS.

C. oblongo-ovata, sangvinea, nitida, supra glabra, subtus pubescens, antennis tibiisque flavo-pallidis, elytris atris, apice sangvineis.

Habitat in Sierra Leona Africæ. D. Afzelius.
2½ lin. longa. Tab. 4. f. 1.

Caput breve, antice truncatum, sangvineum, nitidum, punctis in vertice quibusdam subtilissimis, sulcoque transversali inter oculos profundo impressis. Oculi globosi, prominuli, nigri. Antennæ dimidio corpore longiores, in medio frontis ad latera oculorum insertæ, tenues, filiformes, pallide flavæ, subpubescentes; articulus primus magnus obconicus; 2:us, 3:usque minutæ, nodiformes; 4:us - 10:us subcylindrici, extrorsum non nihil breviores; ultimus acuminatus fuscescens.

Thorax transversus, longitudine duplo latior; antice truncatus, postice vix dilatatus, basi media truncatus, versus angulos antrorum protractus, lateribus subrectis marginatis, supra modice convexus, foveola discoidali utrinque obsolete impressa, sangvineus, nitidissimus, subtilissime punctulatus.

Scutellum triangulare, sangvineum, nitidum, laevissimum.

Elytra thorace quadruplo longiora, eodemque basi dimidio fere latiora, postice parum dilatata, apice & ad humeros obtuse rotundata, abdominis segmentum ultimum non tegentia, angulo apicis interno itidem parum rotundato, supra convexa, humeris parum prominulis, lateribus anguste marginata, subtilissime punctulata, nigra, nitidissima, glabra, apice sanginea.

Corpus subtus sangvineum, nitidum, subtilissime punctulatum, pubescens.

Pedes mediocres, pubescentes; femora sanginea, nitida, subtilissime punctulata; tibiæ tarsiique pallide testacei.

45. CRIOCERIS LIMITATA:

C. oblongo-ovata, pallide testacea, capite, scutello, fasciis maculisque elytrorum, pectore, tibiis & tarsis nigris, fasciis elytrorum impressione limitatis.

Habitat in Brasilia. D. Kreug.

Gall. arcuatæ Oliv. affinis; lin. $3\frac{1}{2}$ longa, Tab. 4. f. 2.

Caput breve, rotundatum, apice truncatum, nigrum, nitidum, inæquale, carinula ante os ele-

vata, atque canalicula inter oculos longitudinali cum impressione transversa crucem formante. Oculi globosi, valde prominuli, nigri. Antennæ dimidio corpore multo longiores, filiformes, basi testaceæ, extrorsum piceo-fuscescentes; articulus primus cylindricus; 2:us minutus, nodiformis; 3:us paullo longior; 4:us - - 10:us elongati, cylindrici, extrorsum sensim fusciores; ultimus acuminatus.

Thorax longitudine dimidio latior, apice truncatus, angustior, mox deinde rotundato-dilatatus, ante angulos posticos subsinuatus, basi in medio truncatus, ad latera antrorsum parum protractus, supra convexus, lateribus margine elevato, flavotestaceus, nitidus, lævis.

Scutellum triangulare, nigrum, nitidum, lævissimum.

Elytra thorace quadruplo longiora, illoque basi dimidio latiora, extrorsum adhuc magis dilatata, apice obtuse rotundata, supra convexa, humeris valde prominentibus, lateribus marginata, subtiliter & crebre punctulata, ultra dimidium testaceo-pallida, posterius pallide flavescentia, fasciis duabus transversis, altera inæquali ante medium, altera vero pone medium sita in maculas duas anguste confluentes divisa, utrisque circumferentia impressa limitatis, nigris; maculæ præterea tres ad basin in lineam dispositæ, quarum duæ humerales, tertia vero ad scutellum communis, punctumque utrinque pone fasciam posteriorem juxta suturam minutum, nigra conspiciuntur.

Corpus subitus pallide testaceum, nitidum, tenuè pubescens, pediore nigro;

Pedes longiusculi, pubescentes, femoribus flava-testaceis, tibiis tarsisque nigris.

46. CRIOCERIS PULCHRA.

G. oblongo-ovata, albido-flavescens, thorace femoribusque latius flavis, capite, medio antennarum, tibiis tarsisque, apice elytrorum, fasciisque tribus transversis nigris.

Habitat in Honduras Americæ Septentrionalis.

D. Weßfelt.

Longitudine $2\frac{1}{2}$ linearum, Tab. 4. f. 3.

Caput subtriangulare, apice truncatum, nigrum, nitidum, vertice levissimo, fronte inter oculos sulcule impressa. Oculi subglobosi, valde prominuli, nigri. Antennæ dimidio corpore multo longiores, filiformes, pubescentes; articulus primus elongatus, obconicus, flavescens; 2:us minutus, nodosus, tertiusque paullo longior, picei; 4:us - 8:us nigri, subcylindrici, æquales; 9:us & 10:us albi; ultimus acuminatus, apice niger, basi albus.

Thorax longitudine paullo latior, antice truncatus, angulis subrectis, lateribus medio rotundatis postice subsinuatis marginatus, flavus totus, supra modice convexus, nitidus, levissimus.

Scutellum triangulare, rufo-piceum, nitidum, levissimum.

Elytra thorace quadruplo longiora eodemque basi dimidio latiora, posterius parum dilatata, apice rotundata, supra modice convexa, humeris sat prominulis, lateribus marginata, subtiliter punctulata, cinereo-albida, nitida, apice fasciisque tribus, quarum prima in ipsa basi sita est, æquali spatio a se

invicem remotis, nec suturam nec marginem attin-
gentibus, nigris.

Pectus & abdomen griseo-flava, subtilissime
punctulata, tenuissimeque pubescentia.

Pedes longiusculi, tenue pubescentes, femori-
bus flavis, tibiis tarsisque nigris.

47. CRIOCERIS DUPLICATA.

C. oblongo-ovata, nitida, pallide flava, capite
thoraceque rufis, elytris basi fasciaque transversa
nigris.

Habitat in Sierra Leona Africæ. D. *Azelius*.

Cr. elatæ *Fabr.* affinis, sed differt capite &
thorace rufis, pectoreque pallido. Lin. $2\frac{1}{2}$ longa.
Tab. 4, f. 4.

Caput rotundatum, ante os truncatum, nitidum,
læve, rufum, ad apicem magis pallescens,
sulco transverso inter oculos impressum. Os pallide
flavum. Oculi rotundati, prominuli, nigri. An-
tennæ dimidio corpore multo longiores, in media
fronte sitæ, tenues, filiformes, pubescentes, pallide
flavæ; articulo primo cæteris longiore, subobconic-
o; 2:o atque 3:o brevioribus; reliquis elongatis
subcylindricis; apicali acuminato.

Thorax longitudine dimidio latior, antice trun-
catus, angulis anticis lateribusque rectis, postice ro-
tundato-subtruncatus, angulis posticis obtusis, su-
pra modice convexus, lateribus tenue marginatis,
subtilissime punctulatus, totus rufus, nitidus, glaber.

Scutellum triangulare, pallide flavum, nitidum,
lævissimum.

Elytra thorace quadruplo longiora, & dimidio latiora, basi humeris rotundatis parum angustata, postice angulo interno utriusque elytri obtuso, ut apice singulatim fere rotundata appareant, supra modice convexa, lateribus anguste marginata, humeris prominulis, subtiliter punctulata, pallide flava, nitida, glabra, basi fasciaque mox pone medium transversa nigris.

Pectus & abdomen subtus pallide flava, subtilissime punctulata, pubescentia.

Pedes mediocres, pallide flavi, pubescentes.

48. CRIOCERIS LITTERATA.

C. pallide flavescentes, ore, antennarum medio pedatoreque fuscis, elytris, sutura baseos signaturisque duabus semicircularibus coeruleoscentibus.

Habitat in Mexico. D. Westfelt.

Cr. fucatæ habitu & magnitudine æqualis, sed colore diversa. Lin. 3 longa. Tab. 4. f. 5.

Caput subrotundum, ante os truncatum, pallide flavescentes, vertice laevi, fronte sulculo inter oculos longitudinali impressa. Os fuscum. Oculi magni, globosi, valde prominentes, nigri. Antennæ dimidio corpore longiores, tenues, filiformes, subpubescentes; articulus primus major, obconicus, secundus brevissimus nodosus, tertiusque secundo paulo longior obconicus, omnes tres pallide flavi; quartus & quintus elongati brunnescentes; 6:us - 8:us æquales, fusi; ultimi tres, quorum apicalis acuminatus, pallide flavi.

Thorax longitudine paulo latior, antice truncatus, angulis rectis, lateribus vix rotundatis mar-

ginatus, basi itidem truncata versus angulos tamen antrorum parum producta, totus pallide flavescentia, supra modice convexus, foveolis discoidalibus binis sat profunde impressis.

Scutellum triangulare, pallide flavum, nitidum, levissimum.

Elytra thorace quadruplo longiora, illoque basi dimidio latiora, posterius dilatata, ad humeros prominulos apiceque rotundata, supra convexa, lateribus marginata, crebre & subtiliter punctulata, pallide flava, sutura a basi ad quartam elytri partem, signaturisque duabus semicircularibus, altera supra humerum sita, litteram C formans, altera interrupta, ex lineis duabus superne versus se invicem flexis, punctoque interjacente, composita, cerasulescentibus.

Pectus fuscum, abdomen pallide flavum, uestrumque pubescens, subnitidum.

Pedes mediocres, pallide flavescentes, pubescentes.

49. CRIOCERIS SEMICINCTA.

C. oblongo-ovata, nigra, capite, thorace, elytris margine apiceque & pedibus testaceo-rufis.

Habitat in Sierra Leonae. D. Afzelius.

Lineas vix 2 longa. Tab. 4. f. 6.

Caput rotundatum, apice truncatum, rufum; nitidum, glabrum, fronte inter antennas sulculo abbreviato impressa. Oculi globosi, prominuli, nigri. Antennae dimidio corpore longiores, tenues, filiformes, rufescentes; articulus primus longior,

obconicus; 2:us & 3:us brevissimi; cæteri subæquales.

Thorax longitudine paullo lator, lateribus parum rotundatis, postice subtruncatus, angulis omnibus subacutis, totus rufus, nitidus, lævissimus, supra valde convexus, anguste marginatus.

Scutellum triangulare, nigrum, nitidum, lævissimum.

Elytra thorace dimidio latiora atque fere quadruplo longiora, apice rotundata, supra convexa, lateribus anguste marginata, humeris parum prominulis, subtiliter punctulato-rugosa, nigra, margine laterali, maculaque apicis lata, juxta suturam ad medium fere elytri antrorum adscendente, sensim evanescente, testaceo-rufis.

Pectus & abdōmen nigra, subtiliter punctulata, tenuē pubescentia.

Pedes mediocres, pubescentes, testaceo-rufi.

50. CRIOCERIS NEGLECTA.

C. oblongo-ovata, testacea, ore, antennisque extorrisum nigris, elytris maculis duabus maximis albis.

Habitat in India Orientali. D. Kymell.

Lin. $1\frac{1}{2}$ longa. Tab. 4. f. 7.

Caput subtriangulare, ante os truncatum, linea transversa inter oculos levius impressa, rufo-testaceum, nitidum, læve. Os nigrum. Oculi globosi, prominuli, nigri. Antennæ dimidio corpore longiores, tenues, filiformes, subpilosæ; articulus primus

primus cæteris longior, obconicus, atque duo insequentes brevissimi, omnes tres testacei; reliqui elongati, cylindrici, nigro-fusci; ultimus apice acutus.

Thorax longitudine duplo latior, antice truncatus, angulis rectis, postice rotundato-subtruncatus, angulis posticis obliquis, lateribus anguste marginatis vix rotundatus, supra modice convexus, rufo-testaceus, nitidus, sublævis.

Scutellum triangulare, rufo-testaceum, nitidum, lævissimum.

Elytra thorace quadruplo longiora atque plus quam dimidio latiora, apice obtuse & singulatim rotundata, supra convexa, lateribus anguste marginata, rufo-testacea, nitida, subtilissime punctulata, macula utrinque magna humerali alteraque adhuc majori, totum fere discum elytri posteriorem occupante, albis.

Corpus subtus rufo-testaceum, pectore paullo obscuriori, nitidum, subtilissime punctulatum, pubescens.

Pedes mediocres testacei, pubescentes; tarsi fusti.

51. CHRYSOMELA GENICULATA. Illiger in litt.

C. gibbosa, supra nigra, subtus nigro-picea, creberrime inæqualiter punctata, femoribus rufis, apice, tibiis tarsisque cœrulescenti-cyaneis.

Trimarcha geniculata, Hoffm. Dejean Catal. d. col. p. 122.

Habitat in Hispania.

C. coriariæ habitu similis, sed minor; lin. 4 longa; mas paullo minor.

Caput rotundatum, ante os truncatum, nigrum, punctatum, fronte depressa, inæqualiter punctato . subrugosa. Oculi laterales, transversi, angustato - oblongi, vix prominuli, brunnei. Antennæ ante oculos insertæ, thorace longiores, moniliformes, crassiusculæ, basi nigro - violaceæ, nitidæ, extrorsum fusco-nigræ, pubescentes; articulus primus magnus, clavatus; 2:us - - 10:us nodiformes, extrorsum paullatim crassiores; ultimus longior, ovato - acuminatus.

Thorax transversus, longitudine duplo latior, antice late & circulariter non vero profunde emarginatus, lateribus ante medium parum dilatatus, posterius non nihil angustatus, basi truncatus, angulis posticis subrectis, supra niger, subnitidus, modice convexus, undique tenue marginatus, subtiliter & crebre punctulatus.

Scutellum longitudine triplo latius, subtriangularē, nigrum, nitidum, laxe.

Elytra magna, thorace fere dimidio latiora & quadruplo longiora, lateribus ovato-dilatata, dupliciter marginata, apice rotundata, supra valde convexa, nigra, nitida, creberrime punctata, punctis inæqualibus, aliis majoribus, aliis minoribus, inter se invicem irregulariter mixtis.

Corpus subtus piceo-nigrum, punctatum, subnitidum.

Pedes longiusculi, validi, nitidi; femora rufa, punctis imprimis ad basin quibusdam impressis, apice cyanea; tibiæ versus apicem incrassatæ, cre-

brius punctatæ, coerulescenti-cyaneæ; tarsi cum tibiis concolores, in mare magis dilatati,

52. **CHRYSOMELA CRUCIGERA.** Schönherr.

C. ovata, nigro-coerulescens, nitida, thorace luteo, macula disci magna antice emarginata, virescenti nigra, elytris luteis, sutura lineaque transversa itidem virescenti-nigris.

Habitat in Brasilia. **D.** Swartz.

C. stauroptera Wied. affinis, sed differt pictura elytrorum, punctis thoracis subtilioribus, atque in elytris haud regulariter seriatis. Lin 4 longa. Tab. 4. f. 8.

Caput subtriangulare, virescenti-nigrum, punctulatum, ante os depresso, fronte sulculo longitudinali impressa. Oculi angustato-oblongi, transversi, laterales, brunnei. Antennæ longitudine dimidi fere corporis, crassiusculæ, nigræ, apice pubescentes; articulus primus crassior, secundus brevior nodosus, atque tertius elongatus, piceo-nigri; 4:us - - 10:us extrorsum gradatim majores; ultimus oblongus, acuminatus.

Thorax transversus, longitudine duplo latior, antice late & profunde emarginatus, angulis anticis valde productis acutis, lateribus tenue marginatis parum rotundatus, basi subbisinuatus, angulis posterioribus rectis, supra convexus, punctis in dorso paucis subtilissimis, in lateribus vero profundioribus atque paullo crebrioribus impressis, luteus, nitidus, macula disci magna, antrorum emarginata, virescenti-nigra.

Scutellum triangulare, apice rotundatum, nigrum, nitidum, laevissimum.

Elytra thorace paullo latiora sed plus quam quadruplo longiora, apice rotundata, supra convexa, punctulata, punctis irregulariter subseriatis, lutea, sutura tota lineaque transversa, pone medium oblique ducta, nec marginem nec suturam attinente, virescenti-nigris.

Pectus & abdomen nigro-coerulescentia, marginibus segmentorum angustissime testaceis.

Pedes mediocres, coerulescentes, nitidi.

53. *CHRYSOMELA SALVIAE*. *Dejean*.

C. ovata, coerulescens, thorace scutelloque dupliciter, elytris vero tripliciter punctulatis, punctis profundissimis elytrorum in series per paria approximatas dispositis.

Habitat in Dalmatia. *D. Dejean*.

C. geminatae valde affinis, sed differt magnitudine majori, colore minus violaceo, scutello punctato, atque imprimis tota superficie superiori oculo acute armato subtilissime & creberrime punctulata. Lin. $3\frac{1}{2}$ longa.

Caput breve, rotundatum, violaceo-coeruleum, subtiliter & parce punctatum, fronte sulculo abbreviato, in angulo linearum duarum ante os profunde impressarum desinente, leviter exarata. Oculi magni, transversi, nigro-brunnei. Antennae thorace paullo longiores, nigræ, basi nitidæ, extrorsum pubescentes, articulo primo crasso, 2:0 - - 10:0 versus apicem sensim crassioribus, ultimo ovato, acuto.

Thorax transversus, longitudine plus quam duplo latior, antice late & profunde emarginatus, angulis anticis antrorum producis, acutis, postice

rotundato - subbisinuatus, angulis fere rectis, lateribus parum rotundatus, anguste marginatus, supra convexus, margine incrassato, violaceo - coerulescens, subtiliter punctatus.

Scutellum rotundato - triangulare, violaceo - coerulescens, nitidum, subtiliter punctatum.

Elytra thorace paullo latiora sed quadruplo fere longiora, apice rotundata, supra convexa, cærulea, punctis crebris minoribus vage adspersis, majoribus vero in series per paria approximatas dispositis. Præter puncta vero jam nominata, superficies capitis, thoracis, scutelli elytrorumque tota, punctulis subtilissimis & creberrimis, nec nisi oculo acute armato conspicuis, quasi reticulato strigosa appetat.

Corpus subtus coerulescens, nitidum, lave.

Pedes mediocres, coerulecentes, nitidi, punctis quibusdam impressis; tarsi subtus ad latera griseo-spongiosi, medio longitudinaliter glabri, nitidi.

54. CHRYSOMELA CIRCUMCINCTA.

C. rotundato - ovata, cærulea, ore, antennarum basi, margineque thoracis & elytrorum testaceo - rufis.

Habitat in Sierra Leona. D. Afzelius.

C. Senegalensi affinis, sed differt corpore magis rotundato, margine thoracis haud incrassato atque punto nigro in lateribus ejusdem nullo. Lin. $2\frac{1}{2}$ longa. Tab. 4. f. 9.

Caput breve, rotundatum, coerulescens, subtilissime punctulatum, ante os foveola semicirculari impressum. Oculi laterales, transversi, brunnei. Antennæ thorace paullo longiores, extrorsum cras.

siores; articulus primus magnus clavatus, & 2:us - 5:us sensim breviores, rufescentes; 6:us - 10:us extrosum gradatim crassiores, & ultimus ovato-acuminatus, nigri, pubescentes.

Thorax basi longitudine plus quam duplo latior, antice angustior, profunde rotundato-emarginatus, lateribus versus basin rotundato-dilatatus, postice parum rotundatus, supra convexus, subtilissime punctulatus, lineola media obsoleta sublaevi, disco viridi-ceruleo, lateribus supra infraque late testaceo-rufis.

Scutellum triangulare, viridi-ceruleo, nitidum, subtilissime punctulatum.

Elytra thorace quintuplo longiora, illoque basi paullo latiora, mox deinde ovato-dilatata, versus apicem rotundata, supra valde convexa, subtiliter & crebre punctulata, cerulea, margine angustius quam in thorace testaceo-rufa.

Pedus & abdomen cerulea, nitida.

Pedes cerascentes, nitidi, femoribus ad basin piceis.

55. EUMOLPUS AZUREUS. Schönherr.

E. azureo-ceruleo, antennarum medio pedibusque cerascentibus, vertice valde gibboso-elevato, thorace subtilissime punctulato.

Habitat in Guinea. D. Colsman.

E. cyaneo affinis, sed differt magnitudine minori, structura vetricis atque thorace subtilius punctulato. Lin. 5. longus.

Caput sub thorace retractum, cyaneum, subtiliter punctulatum; vertex in tuberculum magnum, sulco utrinque profundo ab oculis distinctum, abrupte elevatus. Oculi ovati, laterales, brunnei. Antennæ longitudine dimidii fere corporis, versus apicem compresso-dilatatae; articulus primus crassus, ovatus, & secundus brevis, nigri; 3:us . . 6:us virescentes, obconici; 7:us virescens, 8:us . . 10:us azurei, omnes versus apicem gradatim magis compresso-dilatati, subtilissime punctulati; ultimus maximus, obtuse acuminatus.

Thorax antice angustior, late emarginatus, angulis parum productis, subacutis, posterius sensim dilatatus, basi subtruncatus, undique anguste marginatus, supra valde convexus, lateribus subcompressus, azureo coerulescens, subtilissime punctulatus.

Scutellum rotundato-triangulare, azureo-coerulescens, levissimum, basi foveola punctiformi excavatum.

Elytra thorace dimidio latiora & fere quadruplo longiora, cylindrica, apice rotundata, supra convexa, humeris valde prominulis, tenue marginata, azureo-coerulea, nitidissima, subtiliter vase punctulata; stria suturali ad apicem tenue impressa.

Corpus subtus coerulecenti-azureum, nitidum:

Pedes cyaneo-virescentes, femoribus profunde punctatis, tibiis costato-angulatis.

56. CLYTHRA VARIANS. Sturm.

C. viridis, nitidissima, antennis fuscis, fronte foveolata, thorace lavi, margine elevato punctato-rugoso, elytris subtilissime punctulatis.

Dejean. Catal. d. Coleopt. p. 126.

Var. b. *Coeruleo-virescens.*

Habitat in Brasilia.

Magnitudine *C. quadripunctata æqualis.*

Lin. 4. longa.

Caput retractum, rotundatum, viride, nitidum, subtiliter punctulatum, fovea inter oculos profunde excavata. Oculi oblongi, transversi, parum prominuli, brunnei. Antennæ thoracis basin *vix* attingentes, extrorsum serratæ; articulus primus clavatus, virescens; secundus itidem virescens & tertius fuscus, breves, nodosi; reliqui omnes fusci, compresso-dilatati; ultimus ovatus acuminatus.

Thorax transversus, longitudine duplo latior, antice late emarginatus, basi bisinuatus, lobo versus scutellum retracto obtuse rotundato, angulis omnibus obtusis, supra convexus, lateribus parum rotundatis late, basi vero anguste marginatus, margine laterali elevato reflexo punctato-rugoso, cæterum lævis, æneo-virescens, nitidissimus.

Scutellum elongatum, subtriangulare, convexulum, æneo-virescens, nitidum, læve.

Elytra thorace parum latiora, sed plus quam triplo longiora, cylindrica, apice singulatim rotunda, supra valde convexa, humeris prominulis, æneo-virescentia, nitida, subtilissime punctulata.

Corpus subtus virescens, nitidum, subtilissime punctulatum, pilis griseis obsitum.

Pedes mediocres, virides, nitidi, pilis griseis adspersi.

Var. b. non nisi colore corporis cœruleo-virescenti differt. *Expli-*

Explicatio Tabularum.

Tabula I.

- Fig. 1. Lampyris discoidea*, aucta, mensura longitudinis adjecta.
 — 2. *Copris cornigera*, mas.
 — 3. *Copris cornigera*, femina.
 — 4. *Melolontha æruginosa*.
 — 5. *Rutela Histrio*.
 — 6. *Akis granulifera*.
 — 7. *Helops opacus*.
 — 8. *Tylodes Armadillo*.

Tabula II.

- Fig. 1. Homalinotus deplanatus*.
 — 2. *Amerhinus Ynca*.
 — 3. *Odontoderes Cacicus*.
 — 4. *Lasiopus ciliipes*.
 — 5. *Alocorhinus clericus*, auctus, mensura longitudinis latitudinisque adjecta.
 — 6. *Platyonus nodipennis*.
 — 7. *Phaops Thunbergii*, auctus, addita mensura magnitudinis.
 — 8. *Phaops Albugo*, auctus, adjecta magnitudinis mensura.
 — 9. *Chlorophanus fallax*, auctus, mensura longitudinis latitudinisque adjecta.
 — 10. *Cycloderes catarractus*, auctus, adjecta mensura magnitudinis.
 — 11. *Lixus Sitta*.

Tabula III.

- Fig. 1. Cerambyx zonatus*.

Fig. 2. Dorcadion sericatum.

- 3. *Lema bicruciata*, aucta, adjecta mensura magnitudinis.
- 4. *Casida axillaris*, aucta, mensura magnitudinis adiecta.
- 5. *Casida diaphana*, aucta, mensura magnitudinis adiecta.
- 6. *Casida corrugata*, aucta, mensura magnitudinis adiecta.
- 7. *Galleruca Lanio*.
- 8. 9. 10. *Crioceris mutabilis*, var. a, b & c, auctæ, mensura magnitudinis adiecta.

Tabula IV.

Fig. 1. Crioceris apicalis.

- 2. *Crioceris limitata*.
- 3. *Crioceris pulchra*.
- 4. *Crioceris duplicata*.
- 5. *Crioceris litterata*.
- 6. *Crioceris semicincta*.
- 7. *Crioceris neglecta*.
- 8. *Chrysomela crucigera*.
- 9. *Chrysomela circumcincta*, omnes auctæ, mensura magnitudinis singulorum adiecta.

2.



3.



1.



4.



5.



8.



6.



7.









